

Hepatitis Panel/Acute Hepatitis Panel



CPT: 80074

CMS National Coverage Policy

Medically Supportive ICD Codes are listed on subsequent page(s) of this document.

Coverage Indications, Limitations, and/or Medical Necessity

This panel consists of the following tests:

- Hepatitis A antibody (HAAb), IgM antibody;
- Hepatitis B core antibody (HBcAb), IgM antibody;
- Hepatitis B surface antigen (HBsAg); and
- Hepatitis C antibody.

Hepatitis is an inflammation of the liver resulting from viruses, drugs, toxins, and other etiologies. Viral hepatitis can be due to one of at least five different viruses, designated hepatitis A, B, C, and E. Most cases are caused by hepatitis A virus (HAV), hepatitis B virus (HBV), or hepatitis C virus (HCV).

HAV is the most common cause of hepatitis in children and adolescents in the United States. Prior exposure is indicated by a positive IgG anti-HAV. Acute HAV is diagnosed by IgM anti-HAV, which typically appears within four weeks of exposure, and which disappears within three months of its appearance. IgG anti-HAV is similar in the timing of its appearance, but it persists indefinitely. Its detection indicates prior effective immunization or recovery from infection. Although HAV is spread most commonly by fecal-oral exposure, standard immune globulin may be effective as a prophylaxis.

HBV produces three separate antigens (surface, core, and e (envelope) antigens) when it infects the liver, although only hepatitis B surface antigen (HBsAg) is included as part of this panel. Following exposure, the body normally responds by producing antibodies to each of these antigens; one of which is included in this panel: hepatitis B surface antibody (HBsAb)-IgM antibody. HBsAg is the earlier marker, appearing in serum four to eight weeks after exposure, and typically disappearing within six months after its appearance. If HBsAg remains detectable for greater than six months, this indicates chronic HBV infection. HBcAb, in the form of both IgG and IgM antibodies, are next to appear in serum, typically becoming detectable two to three months following exposure. The IgM antibody gradually declines or disappears entirely one to two years following exposure, but the IgG usually remains detectable for life. Because HBsAg is present for a relatively short period and usually displays a low titer, a negative result does not exclude an HBV diagnosis. HBcAb, on the other hand, rises to a much higher titer and remains elevated for a longer period of time, but a positive result is not diagnostic of acute disease, since it may be the result of a prior infection. The last marker to appear in the course of a typical infection is HBsAb, which appears in serum four to six months following exposure to infected blood or body fluids; in the U.S., sexual transmission accounts for 30% to 60% of new cases of HBV infection.

The diagnosis of acute HBV infection is best established by documentation of positive IgM antibody against the core antigen (HBcAb-IgM) and by identification of a positive hepatitis B surface antigen (HBsAg). The diagnosis of chronic HBV infection is established primarily by identifying a positive hepatitis B surface antigen (HBsAg) and demonstrating positive IgG antibody directed against the core antigen (HBcAb-IgG). Additional tests such as hepatitis B e antigen (HBeAg) and hepatitis B e antibody (HBeAb), the envelope antigen and antibody, are not included in the hepatitis panel, but may be of importance in assessing the infectivity of patients with HBV. Following completion of a HBV vaccination series, HBsAb alone may be used monthly for up to six months, or until a positive result is obtained, to verify an adequate antibody response.

HCV is the most common cause of post-transfusion hepatitis; overall HCV is responsible for 15% to 20% of all cases of acute hepatitis, and is the most common cause of chronic liver disease. The test most commonly used to identify HCV measures HCV antibodies, which appear in blood two to four months after infection. False positive HCV results can occur. For example, a patient with a recent yeast infection may produce a false positive anti-HCV result. For this reason, at present positive results usually are confirmed by a more specific technique. Like HBV, HCV is spread exclusively through exposure to infected blood or body fluids.

This panel of tests is used for differential diagnosis in a patient with symptoms of liver disease or injury. When the time of exposure or the stage of the disease is not known, a patient with continued symptoms of liver disease despite a completely negative hepatitis panel may need a repeat panel approximately two weeks to two months later to exclude the possibility of hepatitis. Once a diagnosis is established, specific tests can be used to monitor the course of the disease.

Indications

1. To detect viral hepatitis infection when there are abnormal liver function test results, with or without signs or symptoms of hepatitis.
2. Prior to and subsequent to liver transplantation.

Limitations

After a hepatitis diagnosis is established, only individual tests are needed.

Visit <https://www.synergylaboratories.com/coverageguidance> to view current limited coverage tests, reference guides, and policy information. To view the complete policy and the full list of medically supportive codes, please refer to the CMS website reference

www.cms.gov/Regulations-and-Guidance/Guidance/Transmittals/Downloads/r17ncd.pdf

Hepatitis Panel/Acute Hepatitis Panel



CPT: 80074

CMS National Coverage Policy

Please refer to the Limitations or Utilization Guidelines section on previous page(s) for frequency information.

The ICD10 codes listed below are the top diagnosis codes currently utilized by ordering physicians for the limited coverage test highlighted above that are also listed as medically supportive under Medicare’s limited coverage policy. **If you are ordering this test for diagnostic reasons that are not covered under Medicare policy, an Advance Beneficiary Notice form is required.**

**Note—Bolded diagnoses below have the highest utilization*

Code	Description
B17.9	Acute viral hepatitis, unspecified
B18.2	Chronic viral hepatitis C
B18.9	Chronic viral hepatitis, unspecified
B19.20	Unspecified viral hepatitis C without hepatic coma
B19.9	Unspecified viral hepatitis without hepatic coma
K74.60	Unspecified cirrhosis of liver
K75.9	Inflammatory liver disease, unspecified
R10.13	Epigastric pain
R10.84	Generalized abdominal pain
R10.9	Unspecified abdominal pain
R16.0	Hepatomegaly, not elsewhere classified
R17	Unspecified jaundice
R53.1	Weakness
R53.81	Other malaise
R53.82	Chronic fatigue, unspecified
R53.83	Other fatigue
R63.4	Abnormal weight loss
R74.01	Elevation of levels of liver transaminase levels
R94.5	Abnormal results of liver function studies
Z01.89	Encounter for other specified special examinations

Visit <https://www.synergylaboratories.com/coverageguidance> to view current limited coverage tests, reference guides, and policy information. To view the complete policy and the full list of medically supportive codes, please refer to the CMS website reference

www.cms.gov/Regulations-and-Guidance/Guidance/Transmittals/Downloads/r17ncd.pdf Last Updated: 1/12/25

Disclaimer:

This diagnosis code reference guide is provided as an aid to physicians and office staff in determining when an ABN (Advance Beneficiary Notice) is necessary. Diagnosis codes must be applicable to the patient’s symptoms or conditions and must be consistent with documentation in the patient’s medical record. The Alliance does not recommend any diagnosis codes and will only submit diagnosis information provided to us by the ordering physician or his/her designated staff. The CPT codes provided are based on AMA guidelines and are for informational purposes only. CPT coding is the sole responsibility of the billing party. Please direct any questions regarding coding to the payer being billed.